

# Non-Variceal Upper GI Bleeding Essential Checklist

To be used in conjunction with Upper GI Bleed guideline, available on microguide.

Name of Doctor :

Bleep :

Patient Escalation Status :

Date and Time :

<p>[ Patient Sticker ]</p> <p>Name</p> <p>DoB</p> <p>Hospital No :</p>
--

	Completed Y/N/NA	Time
<b>RECOGNITION</b>		
Upper GI Bleed diagnosed?		
Aetiologies considered?		
Coagulopathy / Drugs considered? If on aspirin, to continue		
<b>RESUSCITATION</b>		
IV Access obtained? (x2 if possible)		
FBC, UE, LFTs, Group and Save, VBG ?		
Thromboelastogram if complex coagulopathy ?		
<b>PRE ENDOSCOPY MANAGEMENT</b>		
Nil-by-mouth		
Haemoglobin : g/dL	Platelets :	INR: Fibrinogen
Coagulopathy supported?		
Give Tranexemic Acid 1g IV TDS if appropriate.		
Give Omeprazole 40mg IV BD.		
Erythromycin 250mg 30-100 min before endoscopy. (OR if contraindicated, for Metocloperamide 10mg IV)		
<b>If haemodynamically unstable despite optimal resuscitation, consider critical care referral.</b>		
<b>RISK ASSESSMENT</b>		
Glasgow Blatchford Score =		
Senior Reviewed by :		
Endoscopy referral		
<b>POST ENDOSCOPY MANAGEMENT</b>		
Repeat FBC ?		
Post endoscopy treatments ?		
VTE and anticoagulations ?		
Iron replacement, if appropriate ?		