



**Contact Details Name:**

**Hospital**

**Telephone:**

This protocol has 4 pages

## MEDIUM CHAIN FAT OXIDATION DISORDERS – ACUTE DECOMPENSATION

**This protocol covers medium chain acyl CoA dehydrogenase (MCAD) deficiency, HMG CoA synthase deficiency, Carnitine palmitoyl transferase 1 deficiency (CPT1).**

(standard version)

- **Please read carefully. Meticulous treatment is important as there is a high risk of serious complications.**
- **If the instructions do not make sense or a problem is not addressed you must discuss your concerns with the consultant on call.**
- **Intervention should occur whilst the blood glucose is still normal.**

### 1. Background

MCAD deficiency is the most common disorder of fat breakdown and the treatment for some other disorders is similar. For most of the time patients are healthy & do not require a special diet. However infections, fasting, diarrhoea or vomiting can lead to serious illness, with encephalopathy and even sudden death. This results from the accumulation of toxic fatty acids.

The early signs of decompensation may be subtle e.g. lethargy or ‘floppiness’. Always listen to parents carefully as they probably know much more than you do. Hypoglycaemia only occurs at a relatively late stage (or very late) so that blood glucose/BMstix should **not** be relied on. Do not delay treatment just because the blood glucose is not low. The aim should always be to intervene whilst the blood glucose is normal. Treatment aims to prevent mobilisation of fat by providing ample glucose - enterally or intravenously.

### 2. Admission

Most patients who present to hospital will require admission as they are likely to have been having treatment already at home. Only allow the child home if you and the family are entirely happy and you have discussed the problems with the consultant on call. The family must have a clear management plan and be prepared to return if the child does not improve.

- **If there is any doubt at all, the child must be admitted, even if only necessary for a short period of observation.**

### 3. Initial plan and management in hospital

⇒ If the child is shocked or clearly very ill arrange for admission to ITU/High dependency.

⇒ If admitted to metabolic/general ward make a careful clinical assessment including blood pressure and a [Glasgow coma score \(for details click here\)](#), even if the patient does not appear encephalopathic. This allows other staff to recognise if the child deteriorates, particularly around the time of a change of shift.

The following blood tests should be done:

- pH and gases
- Glucose (laboratory and bedside strip test)
- Urea and electrolytes
- Full blood count
- Blood culture

### 4. Management

Management decisions should be based primarily on the **clinical** status. The first decision about therapy is whether the child can be treated orally or will need intravenous therapy.

- Can the child tolerate oral fluids?
- Is the child dehydrated? – Note this can be difficult to assess. The best guide is the difference between the current weight and a recent one when well.

Mild: up to 5% weight loss - may be treated orally but assess carefully.

Moderate or severe: >5% - must be treated with intravenous fluids

- **If there is any doubt at all, put up an intravenous line.**

Treat any infection

#### A. ORAL.

If the child is relatively well and not vomiting, oral feeds may be given.

The emergency regimen should be used. Do not delay. This may be given as regular frequent drinks but if the patient is at risk of vomiting or is nauseated fluid should be given either continuously or as small boluses more frequently. [For more information about the emergency oral management click here](#)

Age (years)	Glucose polymer concentration (g/100ml)	Total daily volume**
0-1	10	150-200 ml/kg
1-2	15	100 ml/kg
2-6	20	1200-1500 ml
6-10	20	1500-2000 ml
>10	25	2000 ml

\* If necessary, seek help from your local dietitian. In an emergency a heaped 5 ml medicine spoon holds approximately 7g of glucose polymer.

\*\* For each drink the volume will generally be this figure divided by 12 and given 2 hourly but if the patient is nauseated or refuses try frequent smaller drinks or a continuous naso-gastric infusion.

Electrolytes should be added to the drinks if vomiting and/or diarrhoea is a problem using standard rehydration mixtures following manufacturer's instructions but substituting glucose polymer solution for water

## **B. INTRAVENOUS.**

If the child is unwell

- Give Glucose 200 mg/kg **at once** (2 ml/kg of 10% glucose or 1ml/kg of 20% glucose) over a few minutes.
- Give normal saline 10 ml/kg as a bolus immediately after the glucose unless the peripheral circulation is poor or the patient is frankly shocked, give 20 ml/kg normal saline instead of the 10 ml/kg.. Repeat the saline bolus if the poor circulation persists as for a shocked non-metabolic patient.
- Continue with glucose 10% at 5 ml/kg/h until next solution ready. – see below
- Quickly calculate the deficit and maintenance and prepare the intravenous fluids
  - Deficit: estimate from clinical signs if no recent weight available
  - Maintenance: Formula for calculating daily maintenance fluid volume (BNF for children) 100ml/kg for 1<sup>st</sup> 10kg then 50 ml/kg for next 10kg then 20ml/kg thereafter, using calculated rehydrated weight. Deduct the fluid already given from the total for the first 24 hours.
  - Give 0.45% saline/10% glucose ([for instructions to make this solution click here](#)).
- Having calculated the deficit and the maintenance, give 1/3 of the total for 24 hours over the next 6 hours and the remainder in 18 hours. If intravenous fluids are still needed, continue with the same solution.
- Recheck the electrolytes every 24 hours if still on intravenous fluids.

- Potassium can be added, if appropriate, once urine flow is normal and the plasma potassium concentration is known.

- Hyperglycaemia can be a problem. If the blood glucose exceeds the 8 mmol/l, start an insulin infusion using the local diabetic protocol rather than reducing the glucose intake. **Strict supervision is essential.**

## **5. Progress:**

**Monitoring:** Reassess after 4-6 hours or earlier if there is any deterioration or no improvement  
Clinical assessment should include [Glasgow coma score \(for details click here\)](#) and blood pressure.

Blood tests: Blood pH and gases  
Glucose (laboratory)  
Urea & electrolytes

⇒ If still obviously encephalopathic continue intravenous fluids but if able to take oral fluids safely, switch to drinks by mouth.

⇒ If deteriorating, seek specialist help without delay.

**6. Re-introduction of oral feeds:** Restart oral feeds as soon as possible; once the child is alert and has stopped vomiting. For more dietary instructions please refer to the to the MCAD dietary guidelines (one for parents and carers and one for dieticians on the BIMDG website) or consult your local dietitian for more details.

**7. Going Home:** Only allow the child home if you and the family are entirely happy and you have discussed the problems with the consultant on call. The family must have a clear management plan and be prepared to return if the child deteriorates.

For further information please refer to:

Saudubray J-M, van den Berghe G, Walter JH. (editors) Inborn Metabolic Diseases. Diagnosis and treatment. 5<sup>th</sup> Edition. Springer 2012

Last reviewed in May 2013