## Appendix J: Moderate Risk (CIWA 10-15) Chlordiazepoxide Alcohol Withdrawal **Protocol Chart**

## Attach to standard inpatient drug chart/patient folder

DRUG (APPROVED		Dose	Additional information	Date						
NAME)										
Chlordiazepoxide		30mg		Day No	1	2	3	4	5	6
Prescribers name	Start date	Route		0800		Χ	Х	Х	Х	Х
		PO		1200		Χ	Х	Х	Х	Χ
Signature	Bleep number	Frequency	Pharmacy	1600		Χ	Х	Х	Х	Х
		variable		2200		Χ	Х	Х	Х	Χ
DRUG (APPR	OVED	Dose	Additional information	Date						
NAME)										
Chlordiazepoxide		20mg		Day No	1	2	3	4	5	6
Prescribers name	Start date	Route		0800	Χ		X	Х	Х	Χ
		PO		1200	Χ		Х	Х	Х	Х
Signature	Bleep number	Frequency	Pharmacy	1600	Χ		Χ	Х	Х	Χ
		variable		2200	Χ		Х	Х	Х	Χ
DRUG (APPROVED		Dose	Additional information	Date						
NAME)										
Chlordiazepoxide		15mg		Day No	1	2	3	4	5	6
Prescribers name	Start date	Route		0800	Χ	Χ		Х	X	X
		PO		1200	X	Χ		Χ	X	Х
Signature	Bleep number	Frequency	Pharmacy	1600	Χ	Χ		Χ	Х	Х
		variable		2200	Χ	Χ		Х	Х	Χ
DRUG (APPROVED NAME)		Dose	Additional information	Date						
Chlordiazepoxide		10mg		Day No	1	2	3	4	5	6
Prescribers name	Start date	Route		0800	Χ	Χ	Х			Х
		PO		1200	Χ	Χ	Х	Х	Х	Χ
Signature	Bleep number	Frequency	Pharmacy	1600	Χ	Χ	Х		Х	Х
		variable		2200	Χ	Χ	Х			
	Pabı	inex <b>2 pai</b>	s IV THREE times	s daily for 2	2 days	s ther	)			•
		lf	no response - dis	continue	_					

Clinical response - 1 pair IV OD for a further 5 days OR until clinical improvement ceases

Please also complete CIWA-Ar scale (overleaf) for appropriate patient monitoring.

## Clinical Institute Withdrawal Assessment of Alcohol (CIWA-Ar) Scale

The CIWA-Ar scale is a validated 10-item assessment tool which quantifies the severity of an individual's alcohol withdrawal symptoms. As such, CIWA-Ar provides guidance on monitoring and benzodiazepine dosage throughout withdrawal.

## Attach to the drug chart and re-check every 4 hours unless CIWA-Ar is worsening, then increase monitoring until patient is stable

Nausea & Vomiting	Tremor					
Ask "Do you feel sick to your stomach? Have you vomited?" Observation.	Arms extended and fingers spread apart. Observation.  0. No tremor					
No nausea and no vomiting     Mild nausea with no vomiting     .     .     .     Intermittent nausea with dry heaves     .     .     Constant nausea, frequent dry heaves and vomiting	Not visible, but can be felt fingertip to fingertip     3.     Moderate, with patient's arms extended     5.     6.     7. Severe, even with arms not extended					
Paroxysmal sweats	Anxiety					
Observation.	Ask "Do you feel nervous?" Observation.					
No sweat visible     Barely perceptible, palms moist     Beads of sweat obvious on forehead     Beads of sweat obvious on forehead     Drenching sweats	No anxiety, at ease     Mildly anxious     Moderately anxious, or guarded, so anxiety is inferred     Equivalent to acute panic states seen in severe delirium or acute schizophrenic reactions.					
Agitation	Tactile disturbances					
Observation.  O. Normal activity  Somewhat more than normal activity  Moderately fidgety and restless  Moderately fidgety and restless  Paces back and forth during most of the interview or constantly thrashes about	Ask "Have you any itching, pins and needles sensations, any burning, any numbness or do you feel bugs crawling under your skin?" Observation.  0. None 1. Very mild itching, pins & needles, burning or numbness 2. Mild itching, pins & needles, burning or numbness 3. Moderate itching, pins & needles, burning or numbness 4. Moderately severe hallucinations 5. Severe hallucinations 6. Extremely severe hallucinations 7. Continuous hallucinations					
Auditory disturbances	Visual disturbances					
Ask "Are you more aware of sounds around you? Are they harsh? Do they frighten you? Are you hearing anything that is disturbing you? Are you hearing things you know are not there?" Observation.  1. Very mild harshness or ability to frighten 2. Mild harshness or ability to frighten 3. Moderate harshness or ability to frighten 4. Moderately severe hallucinations 5. Severe hallucinations 6. Extremely severe hallucinations 7. Continuous hallucinations	Ask "Does he light appear to be too bright? Is colour different? Does it hurt your eyes? Are you seeing anything that is disturbing you? Are you seeing things you know are not there?" Observation.  0. Not present 1. Very mild sensitivity 2. Mild sensitivity 3. Moderate sensitivity 4. Moderate sensitivity 4. Moderately severe hallucinations 5. Severe hallucinations 6. Extremely severe hallucinations 7. Continuous hallucinations					
Headache, fullness in head	Orientation & Clouding of sensorium					
Ask "Does your head feel different? Does it feel like there is a band around your head? Do not rate for dizziness and light-headedness. Otherwise, rate severity.  0. Not present 1. Very mild 2. Mild 3. Moderate	O. Orientated and can do serial additions Cannot do serial additions or is uncertain about the date Disorientated for date by no more than 2 calendar days Disorientated for date by more than 2 calendar days Disorientated for place and/or person					
5. Moderate 4. Moderately severe 5. Severe 6. Very severe 7. Extremely severe	Total CIWA-Ar Score					