Prevention and Management - Coding ICD-11

University Hospitals Sussex

NHS Foundation Trust

New Pathway for Royal Sussex County and Princess Royal Hospitals

There are four different types of MASD

These are:

- 1. Incontinence-Associated Dermatitis, often known as IAD
- 2. Intertriginous Dermatitis, also known as Intertrigo (Intertrigo: where two skin areas may touch or rub together)
- 3. Periwound Maceration
- 4. Peristomal Dermatitis (relating to colostomy, ileostomy/ileal conduit, urostomy, suprapubic catether, or tracheostomy)

1. Incontinence-Associated Dermatitis (IAD) - Coding EK02.22

Actions Required

Complete a Datix and refer for Clinical Images

- ⇒ Keep skin clean and pat dry, avoid rubbing to reduce abrasions to the epidermal layer. Apply a barrier product to restore and protect the skin
- ⇒ Use incontinence shaped pads to absorb urine and faeces. Avoid using the blue flat incontinence sheets/linen savers
- ⇒ Consider the use of appropriate products or devices to divert urine or stool from the skin such as catheters, conveens, and the use of a Flexi-seal™faecal management system

IAD should always be left exposed without dressings

Categorisation	Cleanse	Protect	Images of Product	Examples
Mild Skin Damage	Chose One Method for Cleansing;	Medi Derma-S Total Barrier Cream Every third wash or Twice a day	Total Barrier Com	Processor and the second se
Moderate Skin Damage	Low pH soap such as Senset Foam or Contiplan 7-in-1 Wipes	Cavilon Non Sting Barrier Film Maximum twice per day	Cervion Sector Succession	
Severe Skin Damage		50:50 Ointment Every episode of washing or Fours times per day	LELE LANCENCET 20 March	Servin

Treat areas of **Cutaneous Candidiasis (thrush)** with antifungal treatments

1st Line treatment Clotrimazole 1 % Cream

If using anti-fungal cream for infected area, wait for cream to dry and then apply barrier cream, film or ointment

Senset Foam	150ml	£0.79 each canister	NHS SC Order Code GDW045
Contiplan Wipes	Pack of 25 wipes	£1.91 per pack	NHS SC Order Code VJT188
Dermol 500 Lotion	500 grams	£6.04	Ordered through Pharmacy
Medi Derma-S™ Total Barrier Cream	Individual Sachet 2g (20)	£0.17 each	NHS SC Order Code ELY536
Cavilon™ Non Sting Barrier Film	Foam applicator 1ml (25)	£0.76 each	NHS SC Order Code ELY038
50:50 Ointment	250 grams 500 grams	£1.83 each £3.66 each	Ordered through Pharmacy

If no improvement after 72 hours please consider referral to the Tissue Viability Team for advice and support

2. Intertriginous Dermatitis (Intertrigo) - Coding EK02.20

Examples	Actions Required	
Expiriment	 ⇒ Minimise skin-on-skin contact and friction ⇒ Remove irritants from the skin and protect ⇒ Control or divert the moisture source ⇒ Prevent infection with regular inspection ⇒ Wipes should be left insitu 	

Actions Required

- \Rightarrow Keep skin clean and dry, light patting and not rubbing. This will reduce abrasions to the epidermal layer
- \Rightarrow Allowing the skin between the folds to **completely dry** and be exposed will prevent skin damage
- ⇒ Avoid using dressings and/or talcum powder

Treatment Requirements

- \Rightarrow If skin is broken use **Dermol 500** as a soap substitute
- ⇒ Treat areas of cutaneous yeast and/or fungi infections with topical antifungal treatments such as clotrimazole
- ⇒ Bacterial infections may be treated with topical antibiotics such as Mupirocin ointment
- \Rightarrow **Contiplan wipes** can be placed in to the folds to moisturise and prevent further deterioration

\Rightarrow Avoid using topical steroids

