

What should I avoid?

In the first **48 hours** after injury avoid the following as they might make your symptoms worse:

- Hot baths or showers
- Heat packs or heat rubs
- Drinking alcohol
- Massage

When should I seek help?

- The length of time it takes for your ankle to recover depends on the severity of the original injury. It can take a few months to regain full function.
- If the pain and swelling has not shown improvement within 2 weeks and you are unable to weight-bear please seek further medical advice.

General Advice

- Keep your ankle moving, as pain allows.
- The sooner you commence early exercise, the better your recovery.
- Once you have recovered, continue to keep your body healthy with exercise.
- Strengthen your ankles to prevent future injuries.
- If your symptoms get worse, or do not improve please seek further medical advice.

If you do not understand this leaflet, we can arrange for an interpreter.

إذا كنت لا تستطيع فهم محتويات هذه النشرة فيمكننا عمل الترتيبات لتوفير مترجم شفوي لك.

এই প্রচারপুস্তিকাটি যদি আপনি বুঝতে না পারেন, তবে আপনার জন্য আমরা একজন অনুবাদকের ব্যবস্থা করে দিতে পারি।

如你不明白本單張的內容，我們可安排口譯員服務。

如你不明白本傳單的內容，我們可安排口譯員服務。

اگر مندرجات این جزوه را نمی‌فهمید، ما می‌توانیم مترجم در اختیارتان بگذاریم.

Jeśli masz trudności w zrozumieniu tej ulotki, możemy zorganizować tłumacza.

© Brighton and Sussex University Hospitals NHS Trust

Disclaimer

The information in this leaflet is for guidance purposes only and is in no way intended to replace professional clinical advice by a qualified practitioner.

Ref number: 367.1

Publication Date: August 2013

Review Date: August 2015



Ankle Sprain Advice for Patients



What is an ankle sprain?

- An ankle sprain is one of the most common types of soft tissue injury.
- It occurs when the foot is moved through a greater range of movement than normal. This stretches and weakens the soft tissues and ligaments that support the ankle.

What are the symptoms?

- Pain
- Swelling
- Bruising
- Difficulty moving the ankle normally

How long will it take to recover?

- The initial pain usually settles after a few days and during this time it is important to follow the simple treatment advice in this leaflet to ensure a good recovery.
- The soft tissues (muscles and ligaments) normally take around six weeks to heal, this timeframe may vary as everyone recovers from injuries at different rates.
- You are more likely to sprain your ankle if you have sprained it before. Be careful on uneven surfaces, going up and down stairs, and walking in shoes that are not supportive.

What treatments can I use?

Painkillers

- If your GP has prescribed pain killers, then these should be taken at the recommended dose.

Cold Packs

- A cold pack (ice pack or frozen peas wrapped in a damp towel) can also provide short-term pain relief. Apply to the sore area for up to 15 minutes, every few hours.

Rest

- Try to rest the foot for the first 24-72 hours. However, it is important to maintain flexibility in the ankle so you can gently move it without causing too much pain. This will ensure your ankle does not become too stiff and it will help the healing process.

Elevation

- Raise your ankle above the level of your hips to reduce swelling. You can use pillows to keep the foot up.

Early movement and exercises


- Early weight bearing (putting weight through your injured foot) has been shown to help your ankle heal more quickly. For this reason most people with an ankle sprain will not be given crutches.
- Try to walk as normally as possible as this will help your recovery.

Exercises for your ankle

1. Point your foot up and down.




2. Make circles with your foot. Change direction



3. Place your foot on the floor. Rock your foot from side to side, lifting the inner and then outer border of your foot.



4. Sit with your leg straight out in front of you. Put a towel round your foot and pull it up towards you. Feel a stretch in the back of your calf.



Exercise Illustrations: © PhysioTools Ltd

Please Note: These exercises should not increase your pain. Perform each exercise gently and slowly, only moving as far as feels comfortable. Practice each exercise up to 10 times, 2-3 times a day.