

Bleeding in early pregnancy

You may have experienced some bleeding and/or pain in the early part of your pregnancy which has resulted in you attending the early pregnancy clinic. This leaflet is to give you some information about the bleeding you may have experienced and what happens next.

How common is bleeding in early pregnancy?

Bleeding during pregnancy is relatively common with around one in four women experiencing some bleeding. Around one in five women go on to have a miscarriage. If you have bleeding at any stage during your pregnancy, that causes you concern then you are advised to contact your GP and or midwife for an early assessment.

S/he may refer you to the hospital for an ultrasound scan to establish if the pregnancy is developing as it should. Normal activities of daily living do not cause miscarriage/pregnancy loss and in most cases, bleeding is not a sign of a serious problem. However, it is important (and reassuring) to get assessed by a health professional.

During this time we advise you to use a pantyliner or a sanitary towel instead of a tampon.

When is the most common time to see vaginal bleeding in early pregnancy?

Bleeding /Spotting can be relatively common during the early stages of pregnancy. There could be various reasons.

In early pregnancy you might get some light bleeding when the fetus plants itself into the wall of your womb. This is also known as implantation bleeding, and often happens around the time that your first period after conception would have been due.

There are other less frequent causes of vaginal bleeding which can include bleeding from the neck of the womb (the cervix). Bleeding or spotting may be continuous or intermittent for hours/ days or maybe even weeks. This does not mean that you are miscarrying

or that you will miscarry. Studies have shown that some women who have had bleeding in pregnancy may still go on and have a successful pregnancy.

What investigations do I need if I have bleeding?

If you have bleeding in early pregnancy you may be referred to the Early Pregnancy Unit (EPU) for assessment. This will include a pregnancy test, a scan and possibly blood tests. In the very early stages of pregnancy the scan may need to be done vaginally. This is a safe and (usually) painless examination. Occasionally we may also need to do a vaginal examination.

Do I need any further investigations?

Hopefully the results of all these tests will be reassuring and your pregnancy will continue without any further problems. We often cannot give you a reason why the bleeding happened and we do not normally suggest any further tests.

If the scan does pick up a heartbeat and the baby appears to be the right size according to your dates, this can be very reassuring, even if you are still bleeding.

Research has shown that if you see a heartbeat at 6 weeks of pregnancy, the chances of the pregnancy continuing are 78%.

A heartbeat at 8 weeks increases the chance of a continuing pregnancy to 98% and at 10 weeks that goes up to 99.4%.

So things could still go wrong and sadly sometimes do, but as long as there is a heartbeat, the risk of miscarriage decreases as the weeks go by.

We hope you will find this information helpful and reassuring. The important thing to remember is that whatever you do (having a scan, resting or carrying on with normal activities of daily living), if the bleeding is because the pregnancy is failing, nothing can be done to stop this.

If you have further questions, do not hesitate to ask a member of our team.

Contact numbers

Royal Sussex County Hospital

(Main) 01273 696955
(EPAC) Ext. 4402, Monday-Friday 8-6
(Level 11) Ext. 4013

Princess Royal Hospital

(Main) 01444 441881
(EPAC) Ext. 8370 / Ext. 5686
(Horsted Keynes) Ext. 5685

The Miscarriage Association

17 Wentworth Terrace, Wakefield, WF13QW
tel: 01924 200799 e-mail: info@miscarriageassociation.org.uk
www.miscarriageassociation.org.uk

If you have bleeding and/or pain you can get medical help/advice from:

- Your GP or midwife who may advise you to go to hospital
- Your nearest EPU which can be found at www.earlypregnancy.org.uk/FindUS1.asp add the contact numbers for RSCH AND PRH
- NHS CHOICES 111 when you need medical help fast but it's not a 999 emergency. The service is available 24 hours a day, 365 days a year. Calls are free from landlines and mobile phones.
- BSUH NHS TRUST(2011) GP001 Early Pregnancy Unit Clinical Guidelines.
- Patient UK Information leaflets www.patient.co.uk Women's Health Category
- Royal College of Obstetricians and Gynaecologists www.rcog.co.uk
- National institute for Health and Clinical Excellence: Clinical Guidelines: Nice Pathways www.nice.org.uk

This leaflet is intended for patients receiving care in Brighton & Hove or Haywards Heath

© University Hospitals Sussex NHS Foundation Trust

Disclaimer

The information in this leaflet is for guidance purposes only and is in no way intended to replace professional clinical advice by a qualified practitioner.

Ref number: 392.5
Publication Date: April 2021 Review Date: April 2024

