

Management of ectopic pregnancy

Department of Gynaecology

Summary

What is an Ectopic Pregnancy?

What are the causes of Ectopic Pregnancy?

What are the signs and symptoms?

How is it diagnosed and treated?

How soon can I return to normal activities and work?

How will this affect future pregnancies?

How will it affect me emotionally?


What is an Ectopic Pregnancy?

In normal pregnancy, after the egg has been released from the ovary and fertilized, it travels down the fallopian tube and into the womb (uterus) where it attaches to the lining and continues to develop. In an ectopic pregnancy, the fertilized egg implants outside the womb (usually in the fallopian tube) and continues to grow and enlarge. The fallopian tube is not designed to support a pregnancy and so as the pregnancy grows, the tube is stretched and eventually will rupture.

3 to 5 out of 100 ectopic pregnancies occur in other places such as:

- Cervix or neck of the womb (cervical pregnancy)
- Previous caesarean section scar in the womb (Caesarean scar pregnancy)
- Underdeveloped horn of the womb (cornual pregnancy)
- Ovary (ovarian pregnancy)
- Inside the tummy (abdominal pregnancy)
- In the part of the fallopian tube inside the muscle wall of the womb (Interstitial pregnancy)
- Ectopic pregnancy at any site along with a pregnancy inside the womb (Heterotopic pregnancy).

An ectopic pregnancy can be life threatening because as the pregnancy gets bigger it runs out of space to grow and can potentially rupture(burst) causing severe pain and internal bleeding.



What are the causes of Ectopic Pregnancy?

Ectopic pregnancy occurs in about 1 in 100 pregnancies. The fertilized egg normally spends 4-5 days travelling down the fallopian tube to the womb where it implants. The most common reason for an ectopic pregnancy is damage to the fallopian tube, resulting in a blockage or narrowing which may delay the passage of the fertilized egg allowing it to implant in the tube.

Any woman of child bearing age who is sexual active could be at risk of an ectopic pregnancy.

Your risk increases if:

- You have had a previous ectopic pregnancy.
- You have a damaged fallopian tube, the main causes of damage include: previous surgery to your fallopian tubes including sterilization and previous infections in your fallopian tubes i.e. acute pelvic inflammatory diseases.
- Your pregnancy is a result of an assisted conception: in-vitro fertilisation(IVF) or Intracytoplasmic sperm injection (ICSI).
- You become pregnant on the progesterone only contraceptive pill (mini pill) or when you have an intrauterine device (IUD/Coil).

What are the signs and symptoms?

Each woman is affected differently by an ectopic pregnancy and due to symptoms varying so much, some may have a few symptoms whilst others may have many.

The symptoms include:

- Pain in your lower abdomen- it may be one sided
- Vaginal bleeding
- Pain in the tip of your shoulder – this occurs due to blood leaking into the abdomen
- You may have diarrhoea or pain on opening your bowels
- Severe abdominal pain/lightheadness/fainting/collapse.

How is it diagnosed and treated?

Ectopic pregnancies are sometimes difficult to diagnose. Firstly we will do a urine pregnancy test and if it is positive you will have a scan of your pelvis. This usually will be an internal scan through the vagina. If we cannot confirm a pregnancy in your womb then blood samples will have to be taken to measure your hormone levels. An ectopic pregnancy may be suspected when levels don't rise appropriately. Sometimes we can see the ectopic pregnancy on the scan.

Confirmation of an ectopic pregnancy is usually by a Laparoscopy. This is a 'key-hole' surgical procedure where a telescope is passed through a small incision through your tummy button enabling the doctor to see the womb and fallopian tubes. Once it is proven that there is an ectopic pregnancy, surgery to remove it will be performed.

Surgical treatment

Which is the recommended treatment in this hospital.

This is performed under a general anaesthetic and is the recommended treatment:

- If you are acutely unwell
- There is a live ectopic
- Your hormone level is very high
- The diagnosis is uncertain.

The best operation involves removing the tube affected by the pregnancy, as long as the other tube looks normal. If both tubes look damaged then we aim to just remove the pregnancy and leave the tube intact. Usually you can go home the day after the surgery, depending on your recovery. Occasionally, if the surgery is difficult or complicated an open operation or laparotomy may need to be performed. This is where an incision is made along the bikini line and it may mean a slightly longer stay in hospital.

Medical treatment

In certain clinical cases an ectopic pregnancy can be treated with medication that stops the development of the pregnancy and allows it to be reabsorbed by the body.

This treatment needs to be discussed on an individual basis as it is not suitable for all women.

The treatment involves having an injection called Methotrexate injected into your buttock. (Please refer to separate leaflet explaining all about Methotrexate).

You will need to return to the early pregnancy unit for a series of blood tests to monitor your pregnancy hormones. These blood tests are carried out on day 4, day 7, and day 10 after your treatment. Please be aware that you might experience increased pain or a rise in your blood level on day 4.

About 15% of women need a second injection and a smaller number may need surgery.

Some women have side effects from the treatment, which include increase pain, nausea, mouth ulcers, skin rashes and disturbances to GI System.

How soon can I return to normal activities and work?

Laparoscopic surgery could mean an overnight stay or day case depending on your recovery and time of surgery. We would recommend that you take 2-3 weeks off work and gradually resume your normal activities. Before you leave the hospital, the nursing team will advise you about vaginal bleeding, wound care, infection and resuming sexual intercourse. If you have a laparotomy then your recovery time will be longer and the nursing and medical staff can advise you accordingly.

How will this affect future pregnancies?

Before trying for another baby you should allow yourself time to recover both physically and emotionally. We usually advise you wait until you have your next normal period, but you may find that emotionally you need to wait longer.

If you have had an ectopic pregnancy you have a 10% greater risk of having further ectopics. If your fallopian tube has been removed and the other tube is normal there is still a good chance of conception taking place although it may take a little longer. The most recent research suggests that fertility is reduced by 20-25% (opposed to the expected 50%). This is why we remove the damaged tube, as leaving it behind does not improve your fertility significantly but it does increase your risks of further ectopic pregnancies.

If you have had an ectopic pregnancy you should consult your doctor as soon as you suspect you are pregnant. Early monitoring of the pregnancy is important so that another ectopic pregnancy can be excluded. We are happy to see you for a scan at about seven weeks in the early pregnancy clinic.

When is it ok to start having sex again?

This will depend on how you are feeling after the ectopic pregnancy and what treatment you have had. Medically it is safe to start having sexual intercourse when you have stopped bleeding. We would recommend that you use barrier contraception until after your next period if you are wishing to try and conceive again.

How will it affect me emotionally?

Ectopic pregnancy can be a very traumatic and painful experience. It is likely you would have been rushed to the operating theatre with very little time for psychological adjustment. Much of what has happened to you will have been out of your control, maybe leaving you in a state of shock. You will be recovering from surgery; you and your partner will need to cope with the loss of your baby and often the loss of part of your fertility. Your feelings may vary enormously in the weeks and months following your loss. Listed below are some useful contacts that may help in you and your partner's recovery.

Contact numbers

Royal Sussex County Hospital

(Main) 01273 696955
(EPAC) Ext. 4402, Mon-Fri 8-4
(Day unit) Ext. 7242
(Level 11) Ext. 4013

Princess Royal Hospital

(Main) 01444 441881
(EPAC) Ext. 8370 / Ext. 5686
(Horsted Keynes) Ext. 5685

The Ectopic Pregnancy Trust,

Maternity Unit, The Hillingdon Hospital,
Field Heath Road, Uxbridge, Middlesex, UB8 3NN.
Phone: 01895 238 025 www.ectopic.org.uk

NHS CHOICES 111 when you need medical help fast but it's not a 999 emergency. The service is available 24 hours a day, 365 days a year. Calls are free from landlines and mobile phones.

Further support

Association of Early Pregnancy Units www.earlypregnancy.org.uk

Infertility Network UK www.ectopic.org.uk

Miscarriage Association [www.miscarriageassociation.org.uk/
information/types-of-prgnancy-loss/ectopic pregnancy](http://www.miscarriageassociation.org.uk/information/types-of-prgnancy-loss/ectopic-pregnancy)

- BSUH NHS TRUST(2011) GP001 Early Pregnancy Unit Clinical Guidelines.
- Patient UK Information leaflets www.patient.co.uk
Women's Health Category
- Royal College of Obstetricians and Gynaecologists www.rcog.co.uk
- National institute for Health and Clinical Excellence:
Clinical Guidelines: Nice Pathways www.nice.org.uk



© Brighton and Sussex University Hospitals NHS Trust

Disclaimer
The information in this leaflet is for guidance purposes only and is in no way intended to replace professional clinical advice by a qualified practitioner.

Ref number: 428.4
Publication Date: July 2017
Review Date: July 2019