


Pelvic Organ Prolapse

Department of Gynaecology

Patient Information

What is pelvic organ prolapse?

Pelvic organ prolapse means slipping down of the vagina and/or uterus. Prolapse (slipping down) can affect the front and/or back walls of the vagina and/or the uterus or the vaginal vault (top of the front passage after removal of the uterus).



What causes pelvic organ prolapse?

Pelvic organ prolapse is the result of weakness of the ligaments and muscles that support the vagina and uterus in their normal place. This weakness can result from a number of factors, including pregnancy and child bearing, heavy lifting and straining, going through the menopause as well as inherent weakness of the ligaments and muscles, which can be associated with joint hypermobility. The same applies to stress incontinence of urine, which may co-exist with pelvic organ prolapse.

What are the effects of pelvic organ prolapse?

Not all patients having pelvic organ prolapse have problems. If problems occur, they do not necessarily reflect the degree of prolapse. Some patients have significant bother with minimal degree of prolapse while others have little to note with major degrees of prolapse.

Common symptoms include;

- Sense of bulge or lump down below.
- Dragging sensation, especially towards the end of the day.

- Urinary problems, like stress incontinence (leakage of urine on coughing and sneezing), frequency and /or urgency of micturition as well as difficulty passing urine. Some patients can not pass urine without pushing the prolapse up.
- Bowel symptoms, which may include the need to push prolapse to open the bowel.
- Difficulty with sexual intercourse.

How is this condition diagnosed?

Prolapse is confirmed after an internal examination which is usually carried out while lying on the back and on the left side. In difficult cases, patients might be examined while standing up and sometimes examination under anaesthesia might be necessary to establish the extent of pelvic organ prolapse.

How is the condition treated?

If prolapse is not causing any problems, it is left alone. If it is causing problems, then treatment is offered in the form of a vaginal pessary or surgery.

Vaginal pessaries: These control, rather than cure, prolapse though many patients get symptom relief with them. Commonly used types are the ring, shelf and Gellhorn pessaries. Patients can have sex whilst using a ring pessary but not whilst using a shelf or a Gellhorn pessary. Fitting a pessary for the first time is a matter of trial and error and it can take a few goes to find the right size and shape. They need to be replaced every 6 months to avoid ulceration of the vagina. Pessaries are usually offered to patients who can not have surgery for medical reasons or those who wish to defer surgery for some time, for example to have more children. There is a special information leaflet on the use of vaginal pessaries for pelvic organ prolapse.

Surgery: A number of surgical procedures can be performed to correct pelvic organ prolapse and the choice depends on the type and extent of prolapse as well as your age and future fertility plans. In most cases, a vaginal repair is adequate. The uterus may be removed vaginally at the same time, unless you wish to keep it, for example to have more children, in which case it can be supported strongly to one of the ligaments. Patients with prolapse of the vaginal vault (top of the vagina after

removal of the uterus) may have operations carried out through the vagina, abdomen or laparoscopic to support the top of the front passage. There is a special information leaflet on each of these operations.

Who can I contact with any concerns or questions?

If you have any problems or questions, you can contact:

Princess Royal Hospital, Horsted Keynes Ward:
01444 441881 Ext. 5686

Royal Sussex County Hospital, Level 11:
01273 523191

Urogynaecology Unit at Lewes Victoria Hospital:
01273 474153 Ext. 2178

References / useful link:

<http://www.uptodate.com/contents/pelvic-organ-prolapse-in-women-choosing-a-primary-surgical-procedure>

<http://www.webmd.com/urinary-incontinence-oab/pelvic-organ-prolapse>

<http://www.health24.com/medical/Head2Toe/777-778-783,52704.asp>

<http://www.healthywomen.org/condition/pelvic-organ-prolapse>

<http://emedicine.medscape.com/article/276259-treatment>

<http://pelvicfloor.wordpress.com/2009/12/05/pelvic-organ-prolapse/>

This information has been produced by Dr. Sharif Ismail, Consultant Subspecialist Urogynaecologist.

© Brighton and Sussex University Hospitals NHS Trust

Disclaimer

The information in this leaflet is for guidance purposes only and is in no way intended to replace professional clinical advice by a qualified practitioner.

Reference no.645.3

Publish Date: March 2018 Review Date: March 2020

