

Pressure ulcer prevention

What are pressure ulcers?

Pressure ulcers are areas of damage to the skin and underlying tissue. They are also known as pressure sores or bed sores.


Pressure ulcers may cause pain or lead to an extended hospital stay. They can become infected, leading to blood poisoning or bone infection and in extreme cases can become life-threatening.

What causes pressure ulcers?

Pressure ulcers are caused by a combination of:

- **Pressure** – from body weight pressing against a hard surface, damaging the blood supply to that area.
- **Shearing** – this may occur if you slide down, or are dragged up the bed or chair and may damage the deeper layers of the skin.
- **Friction** – poor moving and handling techniques can remove the top layers of the skin.

Common body sites for pressure ulcers are normally over bone prominence:



Are you at risk of pressure ulcers?

There are things we need to know to help prevent pressure ulcers from developing.

Tell us if you're not eating as much as you used to;

Tell us if you're not moving as much as you used to;

Tell us if you have a sore bottom or heels, hips or elbows;

Tell us if you think there's a problem with your cushion or mattress;

Tell us if you are sleeping in your chair rather than a bed;

Tell us you're having incontinence problems;

Tell us if you have a reduced feeling in your feet;

Tell us if you have had a pressure ulcer in the past;

All the above increase the risk of developing a pressure ulcer and may move you from being '**at risk**' through a '**very high risk**'.



You can change your position without help or prompting.
You have good appetite and no serious health problems.



You may have reduced mobility and require prompting to move regularly – occasional incontinence problems and a poor appetite.



You cannot change your position without help or prompting or you may have reduced feeling to areas of your body or persistent continence problems, poor appetite and general poor health.

Nutrition and hydration

Adequate nutrition and fluids are essential to prevent pressure ulcers. **Please ask for advice if you are concerned.**

What can you and your carer do to help prevent a pressure ulcer?

Regular skin inspections – your skin should be observed for:

- Purple/bluish patches on darker skin
- Red patches on lighter skin, that do not blanch following release of light finger pressure
- Swelling
- Blisters
- Shiny areas
- Dry patches, cracks or calluses.

Report any of these signs to the healthcare team looking after you.

Regular repositioning

If **you can**, change your position **frequently**.

- Please ask staff to assist you to do so, if you have difficulty changing position;
- Make sure that bedding is crease free and not pulled too tightly across you, and that clothing, do not have zips and buttons;
- Make sure shoes/slippers and socks are not too tight as this will restrict your circulation. Please inform nursing staff if you feel any discomfort on this common pressure ulcers sites.

Use of equipment

A mattress or cushion may be provided that will help reduce the amount of pressure on your body. Please notify nursing staff if the equipment provided is not working or is uncomfortable. **Water-filled gloves, synthetic and genuine sheepskins, and doughnut type devices should not be used.** Equipment used for moving or lifting should not be left underneath you.

Inform nursing staff if you use any pressure reducing/relieving equipment in your usual care setting.



Example of the early stage of pressure ulcer development



What happens if you do develop/already have a pressure ulcer?

The Trust is committed to reduce the number of pressure injuries that develop whilst people are in our care. However, unfortunately due to the nature and severity of illness, pressure ulcers may still occur in a small number of people. If this happens appropriate preventive care will be provided and a wound dressing plan will be devised and implemented.

An investigation may also be carried out by the trust wound care specialist. You may require specialist equipment and input from community nurses on discharge.

If you need more information on pressure ulcer prevention and treatment, please ask the nursing staff on the ward and/or request a visit from the Tissue Viability Team.

or visit:

www.your-turn.org.uk

www.stopthepressure.com

Author: Nicola Millard

© Brighton and Sussex University Hospitals NHS Trust

Disclaimer

The information in this leaflet is for guidance purposes only and is in no way intended to replace professional clinical advice by a qualified practitioner.

Ref number: 683.2

Publication Date: May 2019 Review Date: May 2021

